



Investigation on *in-vitro* Antidiabetic and Hepatoprotective activity of the leaves of *Bauhinia phenicea* Wight & Arn, Fabaceae

Sonia Johny*, Indira G, Vijishna L V

College of Pharmaceutical Sciences, Govt. Medical College, Thiruvananthapuram - 695011, Kerala, India.

ARTICLE HISTORY

Received: 22.07.2021

Accepted: 30.08.2021

Available online: 30.09.2021

DOI:

10.5530/ajphs.2021.11.17

Keywords:

Bauhinia phenicea, Hepatoprotective, MTT assay, Antidiabetic, α -Amylase inhibition

*Corresponding author:

Phone : + 91 - 7034244150

Email : soniajohny95@gmail.com

ABSTRACT

Bauhinia phenicea Wight & Arn. (Syn: *Bauhinia benthamii* Bedd.) belonging to the family: Fabaceae is an ornamental medicinal climbing shrub endemic to Western Ghats. The plant parts have been used traditionally for treating wounds, diabetes, skin allergies, fungal infections and worm disturbances. The Pharmacognostical and Preliminary phytochemical studies have been reported on the leaves. The plant contains steroids, terpenoids, phenols, tannins, saponins, alkaloids and flavonoids as phytoconstituents. No detailed reports are available on the hepatoprotective and antidiabetic activity of the leaves, hence attempt was undertaken to investigate the hepatoprotective and antidiabetic activity on the leaves of *Bauhinia phenicea* by *in-vitro* methods like MTT assay and α -Amylase inhibition assay respectively. The study revealed that the leaf possess significant antidiabetic and hepatoprotective activities which justifies the therapeutic potential and traditional uses of the plant.

INTRODUCTION

Healing with medicinal plants is holistic and as old as mankind.⁽¹⁾ Enormous number of alkaloids, glycosides, antibiotics, and other phytoconstituents have been isolated from plants, identified and used as curative agents. The Genus *Bauhinia* is a large and diverse tropical and subtropical genus comprising of approximately 300 species, belonging to the family leguminosae or fabaceae. Most of them possessed bilobed leaves and has been the subject of a number of regional treatments. Species of *Bauhinia* display a wide range of morphological variations and habit type, being found as trees, shrub, herbs or lianas.⁽²⁾

α -amylase is one of the key digestive enzyme which catalyses hydrolysis of starch to maltose in the beginning and at the end convert it to glucose therefore by inhibiting the activity of α -amylase. One can think about the control of diabetes as diabetes is related to increase in glucose level inside body and α -amylase is responsible for the conversion of starch into many simple sugars like dextrin, maltose and glucose.⁽³⁾ The activity of α -amylase is

correlated to the elevated postprandial hyperglycemia therefore control of this enzyme activity becomes significant in the treatment of diabetes.⁽⁴⁾

Oxidative stress becomes a detrimental condition when (Reactive oxygen species) ROS is excess. ROS could damage biological molecules such as proteins, deoxynucleic acid, and lipid membranes, which cause apoptotic or necrotic cell death by disrupting cellular function and integrity⁽⁵⁾. This harmful oxidative stress has been considered as a major cause of cell injuries, which leads to many diseases such as Alzheimer's, cancer, and cardiovascular diseases⁽⁶⁾. ROS generated from mitochondria and other sources causes cell damage through mechanisms involving lipid peroxidation and subsequent liver injury. This also activates many kinases to catalyse degenerative progression in liver tissues.⁽⁷⁾ H_2O_2 which occurs as a byproduct of oxidative stress has been implicated in triggering apoptosis in various cell types to evaluate hepatoprotective activity.⁽⁸⁾ Herbal products which consist of many compounds have been used by the people and which has become the dependable source of

remedy for various diseases. Hence, the discovery of new potential drugs from medicinal plants with fewer side effects for the treatment of diabetes and liver diseases has to be continued.

Bauhinia phenicea Wight & Arn is an ornamental medicinal shrub endemic to Western Ghats which is mainly found in the states of Karnataka, Kerala and Tamil Nadu.⁽⁹⁾ The plant, known as vallimantharam in Malayalam, Kembu yathaballi in Kannada, Thukarakkali in Tamil, Raktakanchana or Ashmantaka in Sanskrit is a large climbing shrub having a height of 40- 45 feet, with alternate, bilobed leaf and scarlet red coloured flowers. *Bauhinia phenicea*, with common names Crimson mountain ebony or scarlet Bauhinia consists of dark brown, hard, deeply fissured bark and widely spread crown. Traditional claims reveal that the plant is useful for wound healing, diabetes, skin allergies, fungal infections and worm disturbances.⁽⁹⁾ The antioxidant, anticancer⁽¹⁰⁾ and antimicrobial⁽¹¹⁾ activity of *Bauhinia phenicea* leaves has been reported. Hence this study was designed to carry out investigation on the hepatoprotective and antidiabetic activities of the chloroform extract of the leaves of *Bauhinia phenicea* Wight & Arn. by MTT assay of H₂O₂ induced peroxidation on HepG₂ cells and α -amylase inhibition by starch iodine method respectively.

MATERIALS AND METHODS

Collection of the Plant Material

The plant material was collected from Karyavattom, Trivandrum District, and Kerala, India in September 2020. The plant was authenticated by Dr. T.S. Swapna, Department of Botany, Karyavattom Campus. A voucher specimen (Voucher No. KUBH 10731) has been deposited for future reference. The leaves were separated from other plant parts, washed, cleaned, dried and powdered.

Reagents

All the reagents used were of analytical grade obtained from Universal chemicals and scientific industries, Kerala

Preparation of Extracts⁽¹²⁾

50g of dried coarse powder of leaves packed in a soxhlet-extractor and extracted successively with solvents like petroleum ether (60-80°C), chloroform, ethyl acetate and methanol. The extraction was continued each times till the solvent in the thimble of extractor become clear. The marc obtained after the methanol extraction was dried and macerated with chloroform water for 24 hrs and filtered to obtain the water extract. All the extracts

Table 1 : Mean absorbance and percentage inhibition of Chloroform extract of *Bauhinia phenicea* Wight & Arn. and standard Acarbose by Starch Iodine assay.

Sample	Concentration µg/ml	Mean Absorbance	Percentage inhibition
Control	-	0.313 ± 0.0002	-
Chloroform extract	25	0.395 ± 0.001	27.23
	50	0.432 ± 0.0002	38.21
	100	0.514 ± 0.0009	63.901
	200	0.554 ± 0.0001	76.93
Acarbose	25	0.415 ± 0.002	31.63
	50	0.474 ± 0.001	51.95
	100	0.538 ± 0.001	71.73
	200	0.563 ± 0.0008	80.32

Values are expressed in Mean ± SEM, n = 3, The value were compared with control using Dunnetts multiple comparison and P value obtained was less than 0.0001, *** = p < 0.0001. Therefore, considered as significant

collected were evaporated in a water bath and the dried extracts were stored in a desiccator till required. All the extracts were evaluated for in-vitro anti-oxidant activity. Chloroform extract showed good antioxidant activity hence was selected for carrying out in-vitro antidiabetic and hepatoprotective activity.

Hepatoprotective activity analysis by MTT assay^(13,14,15,16,17)

The hepatoprotective activity of the test samples were evaluated using HepG2 cell line. IC₅₀ concentration of H₂O₂ (30 µg) was used as a cytotoxic agent. Confluent HepG2 cells were cultured in growth media (DMEM + 10% FBS) at a density of 5 × 10⁴ cells/well in a 96-well tissue culture plate and incubated overnight. Post incubation, cells were treated with varying concentrations of the test sample (25, 50, 100 µL) and incubated for 24 h, thereafter; IC₅₀ concentration of H₂O₂ (30 µg) was added and allowed for further 24 h incubation. Post incubation, the treated cells were washed with PBS and incubated with MTT containing growth media. Finally, the medium was removed, and the formazan crystals were dissolved using 150 µl DMSO (100%). The optical density was measured at 570 nm. Untreated cells were kept as control and percentage cell viability in treated cells were calculated.

Antidiabetic activity by α-amylase inhibition by starch iodine method^(3,4,18)

The assay mixture was about 120 µl of 0.02M sodium phosphate buffer (pH 6.9), 1.5 ml of α-amylase and plant extracts at a concentration from 25-200 µg/ml (w/v) was incubated at 37°C for 10 min. After that, soluble 1% starch was added at each reaction mixture and incubated at 37 °C for 15 min. Then 60 µl of 1 M HCl was added to the reaction mixture to stop the enzymatic reaction and immediately 300 µl of iodine reagents was added. If any change in colour was noted and at 620 nm the absorbance was read.

Statistical analysis

Statistical analysis was done by using one-way ANOVA followed by Dunnet's test. P values lesser than 0.05 were considered as significant using Graph Pad Prism software version 9.


RESULTS

The percentage inhibition of α-amylase enzyme was found to increase with increase in concentration of chloroform extract of leaves of *Bauhinia phenicea* Wight & Arn. The values are tabulated in Table no.1. IC₅₀ values of standard Acarbose and chloroform extracts were compared and is depicted in Graph no. 1.

The protective effect of the chloroform extract on cell death induced by oxidation of H₂O₂ was evaluated. The IC₅₀ of H₂O₂ was investigated by inducing oxidation on HepG2 cells whereby percentage cell viability decreased with increasing concentration of H₂O₂. The values are depicted in Table no. 2 and IC₅₀ was found to be 30.08 µM.

IC₅₀ of H₂O₂ was found to be 30.08 µM. The cell viability of HepG2 cells was found to increase with increase in concentration of the leaf extract i.e, in a dose dependant manner and the obtained values have been tabulated in Table no.3.

Values are expressed in mean ± SEM, n=3. The value were compared with control using Dunnet's multiple comparison and P value obtained was less than 0.05, **= p <0.05. Therefore,



Graph 1 : Comparative IC₅₀ (µg/ml) value Chloroform extract of *Bauhinia phenicea* Wight & Arn. and standard Acarbose

Table 2 : MTT result of Hydrogen Peroxide.

Concentration (µM)	Percentage cell viability
6.25	82.15
12.5	73.52
25	61.36
50	47.15
100	36.59

considered as significant.


DISCUSSION

Plant drugs are of great value in the field of treatment and cure of diseases. The preliminary phytochemical evaluation of chloroform extract of the leaves of *Bauhinia phenicea* revealed the presence of large number of secondary metabolites such as flavonoids, saponins and steroids. These secondary metabolites might be responsible for traditional claims and therapeutic activity of this plant. From different literatures, it was reported that the plant has antioxidant, antimicrobial and anticancer activity.

The percentage inhibition of alpha amylase enzyme was found to increase with increase in concentration of chloroform leaf extract. The activity of α-amylase is correlated to the elevated postprandial hyperglycemic conditions. Inhibition of α-amylase enzyme decreases the conversion of carbohydrates into glucose in the blood thereby reducing the chance for hyperglycemia. Thus the chloroform leaf extract of *Bauhinia phenicea* Wight & Arn inhibited α-amylase enzyme thereby substantiating its antidiabetic activity and confirming the traditional claim of the plant

Table 3 : Percentage cell viability when treated with chloroform extract of *Bauhinia phenicea* Wight & Arn.

Concentration	Absorbance	Percentage Cell Viability
Control	0.415	50
H ₂ O ₂ (30 µM)+ 25µg/ml chloroform extract	0.422	55.82
H ₂ O ₂ (30 µM)+ 50 µg/ml chloroform extract	0.490	64.81**
H ₂ O ₂ (30 µM)+ 100 µg/ml chloroform extract	0.583	77.16**

**Fig 1 :** Percentage cell viability during evaluation of Hepatoprotective activity of chloroform extract of *Bauhinia phenicea* Wight & Arn. at different concentrations

HepG2 cells (Human liver carcinoma cells) have been exploited as a model to study the hepatoprotective effect of the test compound in this study. H₂O₂ is also widely used as an inducer of oxidative stress in in-vitro models, which could lead to cell death⁽¹⁹⁾. In addition, it was reported that the cell damage effect induced by H₂O₂ could be attenuated by treating with antioxidants⁽²⁰⁾. Therefore, the protective effect of the test compounds on cell death induced by H₂O₂ was investigated by HepG2 cell model. The percentage cell viability of HepG2 cells was found to increase with increase in concentration of the leaf extract in a dose dependant manner. This may be due to the protective effect induced over the H₂O₂ treated HepG2 cells, by the flavonoids or antioxidant principles present in the chloroform extract of *Bauhinia phenicea*.

CONCLUSION

Diabetes and liver damage are among the major health problems worldwide which may even lead to death. Continuous research for effective natural medicine with lesser side effects is hence a requirement of today's world. The present study showed that the chloroform leaf extract of *Bauhinia phenicea* Wight & Arn possess significant antidiabetic and hepatoprotective activity in a dose dependant manner. The results of this study directs further researches to evaluate the therapeutic potentialities of the phytoconstituents present in the leaf of *Bauhinia phenicea* Wight & Arn. Further *in-vitro* and *in-vivo* studies are to be conducted for the substantiation of the traditional claims of the drug thereby changing this plant into a future promising drug.

ACKNOWLEDGEMENT

The authors are grateful to the Head of the Department Dr. Joyamma Varkey of College of Pharmaceutical Sciences, Government Medical College, Trivandrum and Mr. Austin P, Managing Director of Atmic Biotech Solutions, Trivandrum for providing the facilities for this work.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Biljana Bauer Petrovska. Historical review of medicinal plants' usage. *Pharmacognosy Reviews*.2012;6(11):1-5
2. V. Vadivel, H.K. Biesalski. Nuts and seeds in health and disease prevention. 2011:941-949
3. Alexander R. Maltodextrins: production, properties and applications. In: Schenk F, Hebeda R. ed. Starch hydrolysis products; Worldwide technology: production and applications, New York; 1992. 62122.
4. Eichler HG, Korn A, Gasic S, Prison W, Businger J. The effect of new specific α -amylase inhibitor on post-prandial glucose and insulin excursions in normal subjects and type 2 (non-insulin dependent) diabetic patients. *Diabetologia* 1984;26 Suppl 4:278-81.
5. Valko, M., Leibfritz, D., Moncol, J., Cronin, M.T., Mazur, M., and Telser, J., "Free radicals and antioxidants in normal physiological functions and human disease". *The International Journal of Biochemistry & Cell Biology*. 2007; 39(1):44-84.
6. Markesbery, W.R., "Oxidative stress hypothesis in Alzheimer's disease". *Free Radical Biology & Medicine*. 1997. 23(1). 134-147.
7. Hongtan Wu et al. Hepatoprotective effect of Polyphenol enriched fraction from Folium microcos on oxidative stress and apoptosis in Acetaminophen induced liver injury in mice. *Oxidative Medicine and Cellular longevity*. 2017;1(1): 1-15
8. Hadi Gaffari, Behrouz Jalali Ghassam, HS Prakash. Hepatoprotective and cytoprotective properties of Hyptis suaveolens against oxidative stress- induced damage by CCl₄ and H₂O₂. *Asian Pacific Journal of Tropical Medicine*. 2012; 1(1): 868-874
9. Alby Alphons Baby, Regi Raphael K. First step towards unraveling the medicinal properties of an endemic traditional medicine, *Bauhinia phenicea* Wight & Arn bark. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2015; 7(9): 403- 405
10. Alby Alphons Baby, Regi Raphael K. Analysis of antioxidant property, cytotoxicity and anti-tumour efficiency of *Bauhinia phenicea*. *Asian Journal of Pharmaceutical And Clinical Research*. 2018; 11(12): 277-279
11. Alby Alphons Baby, Regi Raphael K. Evaluation of the folk claim and identification of pharmacological active principles in *Bauhinia phenicea* leaves. *Asian Journal Of Pharmaceutical And Clinical Research*. 2016; 9(4): 1-4
12. Kokate.C.K. Practical Pharmacognosy, 4th edition, Vallabhprakashan, Delhi,1992; 14-23,115-121.
13. Zhong K, Li XJ, Gou AN, Huang YN, Bu Q, Gao H. Antioxidant and cytoprotective activities of flavonoid glycosides-rich extract from the leaves of *Zanthoxylum bungeanum*. *Journal of Food Nutrition and Research*. 2014; 2(7):349-56.
14. *In vitro* efficacy and safety of poly-herbal formulations. Chandrasekaran CV, Sundarajan K, David K, Agarwal A Toxicol In Vitro. 2010 Apr; 24(3):885-97.
15. Siddiqui MA, Ali Z, Chittiboyina AG, Khan IA. Hepatoprotective Effect of Steroidal Glycosides From *Dioscorea villosa* on Hydrogen Peroxide-Induced Hepatotoxicity in HepG2 Cells. *Frontiers in Pharmacology*. 2018; 9(1):797.
16. Parimal P. Katolka, Nandakishore J. Duragkar: Hepatoprotective activity of stem bark of *Bauhinia tomentosa*. *International Journal of Researches In Biosciences, Agriculture & Technology*. 2014; 2(2): 219-225
17. Yahya F, Mamat S S, Kamarolzaman M F et al. Hepatoprotective Activity of Methanolic Extract of *Bauhinia purpurea* Leaves against Paracetamol-Induced Hepatic Damage in Rats. *Evidence Based Complementary & Alternative Medicine*. 2013;1(1): 1-10
18. Priyanka Jaiswal, Padma Kumar. Alpha Amylase inhibitory activity of different extract of bark of *Albizia lebbeck* (L.) Benth. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2017; 9(8): 119-122
19. Zhang, J., Melton, L.D., Adaim, A., and Skinner, M.A., "Cytoprotective effects of polyphenolics on H₂O₂-induced cell death in SH-SY5Y cells in relation to their antioxidant activities". *European Food Research & Technology*.2008; 228(1): 123-131.
20. Jia, Y., Ji, L., Zhang, S., Xu, L., Yin, L., Li, L., Zhao, Y., and Peng, J., "Total flavonoids from *Rosa Laevigata* Michx fruit attenuates hydrogen peroxide induced injury in human umbilical vein endothelial cells". *Food & Chemical Toxicology*. 2012; 50(1): 3133-3141.



Cite this article : Sonia Johny, Indira G, Vijishna L V
Investigation on *in-vitro* Antidiabetic and Hepatoprotective activity of the leaves of *Bauhinia phenicea* Wight & Arn, Fabaceae
Asian J. Pharm. Hea. Sci.. 2021;11(3):2518-2522. DOI : 10.5530/ajphs.2021.11.17